

SOME MEDICATIONS NEED TO BE STOPPED (as much as 2 weeks) PRIOR TO SURGERY

Make your physician aware if you are taking any
medication that contains

ASPIRIN or any medication on this list

| | | |
|----------------|-----------------------|------------------------|
| Alka-Seltzer | Aspirin | Arthritis Pain Formula |
| Bayer Products | Bufferin | Cortisone Medication |
| Excedrin | Fiorinal | IBUPROFEN |
| Midol | Pepto-Pismol | MOTRIN |
| Percodan | Propoxyphene Compound | Sinutab |
| Soma Compound | St. Joseph Aspirin | Triaminicin |
| Trigesic | Advil | Aleve |
| Anaprox | Motrin | Naprosyn |
| Nuprin | Plavix | Actifed |
| Ansaid | Arthritis Bufferin | Asriptin |
| BC Tablets | Coumadin | Doxycycline |
| | Entab-650 | Goody's Powder |
| Heparin | Hydrocortisone | Lovenox Injection |
| Nalproxyn | PanwarfinSparine | Phentermine |
| Predisone | Relafen | Salicylates |
| Sinex | Vitamin E | Warfaren |

I have received a copy of the Hickory Plastic and Reconstructive Surgery Center

Medicines To Be Stopped Prior To Surgery

And agree to discuss any of these medications I am taking with my physician.

Patient Signature

Date

Print Patient Name

Chart No.